

INDEX

COPIEA 1984, NOS. 1-4

NEW NAMES

- Allenichthys* 27-28, 30, 38, 42
Barathronus multidens 579-584
Barathronus unicolor 579-584
Bulbonaricus brucei 565-571
Cichlasoma socolofi 933-940
Cyprinodon verecundus 58-68
Echiodon anchipterus 410-422
Echiodon coheni 410-422
Emoia rufilabialis 571-578
Emoia taumakoensis 571-578
Gorgasia barnesi 404-409
Kuitterichthys 27-28, 32, 37-38, 41-42
- Lumiconger arafura* 72-81
Nanachromis transvestitus 82-86
Pavorajini 45, 52, 55
Phyllophryne 27, 29, 31, 39-42
Physalaemus coloradorum 902-909, 911-912, 914-920
Plectrohyla pokomchi 390-397
Plectrohyla tecunumani 390-397
Rana yavapaiensis 940-948
Rana zweifeli 398-403
Rhycherus gloveri 68-72
Texasophis wilsoni 660-661
Urophycis musicki 1008-1009

NEW AND ESTABLISHED NAMES

- Abantennarius* 34
Abramis 3, 11-12, 265
Acacia spp. 854
Acanthochromis polyacanthus 190, 629, 637
Acanthopterygii 4, 9-12
Acanthuridae, 14, 190
Acanthurus
 dussumieri 14
 mata 14
 olivaceus 14
 triostegus 758-761
 xanthopterus 14
Acari 755
Acentrophorus 499
Acer rubrum 976
Acipenser 476, 481, 497
Acridoheres tristis 268
Acris crepitans 289
Acrochordus 641
Acropora spp. 192
Adinia xenica 797-800
Adrianichthyidae 833
Aeduella 497
Aethedontus 497
Agama agama 267-274
Agave sisalana 748
Agkistrodon
 bilineatus 771
 contortrix 773
 controrix mokeson 772
 piscivorus 772-774
Ahaetulla prasina 642, 647
Alectina 158
Alestes 263
 longipinnis 263
 spp. 263
Allenichthys 27-28, 30, 38, 42
 glauerti 28, 38
Alligator mississippiensis 153-154, 160, 839-843
Allondichthys hubbsi 235, 238
Alosa
- pseudoharengus* 878-882
sp. 205-206, 208
- Alsophiini* 645
- Ambassis* 12 *ranga* 4
Ambloplites rupestris 151
Ambrosia 813 *dumosa* 812
Amblyglyphidodon aureus 198
Amblygobius
 decessatus 189
 phaelena 189-190
Amblyodipsas 644 *polylepis* 642, 644
Amblyraja 52
Ambystoma 161-169, 299, 324, 327, 681, 739, 741,
 745, 866
 dumerilii 742
 gracile 672-683, 863
 jeffersonianum 161-168, 324, 333-335
 jeffersonianum × *Ambystoma texanum* 335
 laterale 324, 333-334
 laterale × *Ambystoma texanum* 335
 macrodactylum 863
 maculatum 161-168, 324, 326, 331-335, 739-740,
 742, 745, 866, 869
 mexicanum 740, 742-743
 opacum 167, 744
 ordinarium 866-867
 platineum 324-337
 platineum × *Ambystoma maculatum* 334-335
 platineum × *Ambystoma texanum* 324, 328, 334-335
 rosaceum 866-867
 texanum 924-937, 739-742, 744-745, 862-868
 texanum × *Ambystoma jeffersonianum* 335
 texanum × *Ambystoma laterale* 334
 tigrinum 740-742, 866
 tremblayi 324-325, 329-330, 333-335
Ambystomatidae 739-747
- Amia* 476, 479-482, 484-485, 490-499, 649
Amiurus 97, 100 *lupus* 97
Ammocrypta beani 388
Ammodytes 11-12

INDEX

COPIEA 1984, NOS. 1-4

NEW NAMES

- Allenichthys* 27-28, 30, 38, 42
Barathronus multidens 579-584
Barathronus unicolor 579-584
Bulbonaricus brucei 565-571
Cichlasoma socolofi 933-940
Cyprinodon verecundus 58-68
Echiodon anchipterus 410-422
Echiodon coheni 410-422
Emoia rufilabialis 571-578
Emoia taumakoensis 571-578
Gorgasia barnesi 404-409
Kuitterichthys 27-28, 32, 37-38, 41-42
- Lumiconger arafura* 72-81
Nanachromis transvestitus 82-86
Pavorajini 45, 52, 55
Phyllophryne 27, 29, 31, 39-42
Physalaemus coloradorum 902-909, 911-912, 914-920
Plectrohyla pokomchi 390-397
Plectrohyla tecunumani 390-397
Rana yavapaiensis 940-948
Rana zweifeli 398-403
Rhycherus gloveri 68-72
Texasophis wilsoni 660-661
Urophycis musicki 1008-1009

NEW AND ESTABLISHED NAMES

- Abantennarius* 34
Abramis 3, 11-12, 265
Acacia spp. 854
Acanthochromis polyacanthus 190, 629, 637
Acanthopterygii 4, 9-12
Acanthuridae, 14, 190
Acanthurus
 dussumieri 14
 mata 14
 olivaceus 14
 triostegus 758-761
 xanthopterus 14
Acari 755
Acentrophorus 499
Acer rubrum 976
Acipenser 476, 481, 497
Acridoheres tristis 268
Acris crepitans 289
Acrochordus 641
Acropora spp. 192
Adinia xenica 797-800
Adrianichthyidae 833
Aeduella 497
Aethedontus 497
Agama agama 267-274
Agave sisalana 748
Agkistrodon
 bilineatus 771
 contortrix 773
 controrix mokeson 772
 piscivorus 772-774
Ahaetulla prasina 642, 647
Alectina 158
Alestes 263
 longipinnis 263
 spp. 263
Allenichthys 27-28, 30, 38, 42
 glauerti 28, 38
Alligator mississippiensis 153-154, 160, 839-843
Allondichthys hubbsi 235, 238
Alosa
- pseudoharengus* 878-882
sp. 205-206, 208
- Alsophiini* 645
- Ambassis* 12 *ranga* 4
Ambloplites rupestris 151
Ambrosia 813 *dumosa* 812
Amblyglypidodon aureus 198
Amblygobius
 decessatus 189
 phaelena 189-190
Amblyodipsas 644 *polylepis* 642, 644
Amblyraja 52
Ambystoma 161-169, 299, 324, 327, 681, 739, 741,
 745, 866
 dumerilii 742
 gracile 672-683, 863
 jeffersonianum 161-168, 324, 333-335
 jeffersonianum × *Ambystoma texanum* 335
 laterale 324, 333-334
 laterale × *Ambystoma texanum* 335
 macrodactylum 863
 maculatum 161-168, 324, 326, 331-335, 739-740,
 742, 745, 866, 869
 mexicanum 740, 742-743
 opacum 167, 744
 ordinarium 866-867
 platineum 324-337
 platineum × *Ambystoma maculatum* 334-335
 platineum × *Ambystoma texanum* 324, 328, 334-335
 rosaceum 866-867
 texanum 924-937, 739-742, 744-745, 862-868
 texanum × *Ambystoma jeffersonianum* 335
 texanum × *Ambystoma laterale* 334
 tigrinum 740-742, 866
 tremblayi 324-325, 329-330, 333-335
Ambystomatidae 739-747
- Amia* 476, 479-482, 484-485, 490-499, 649
Amiurus 97, 100 *lupus* 97
Ammocrypta beani 388
Ammodytes 11-12

- dubius* 243
hexapterus 242–244, 1013
marinus 242–243
personatus 243–244
tobianus 4
- Ammodytidae 1012–1014
- Amphiacusta sp. 219
- Amphibia 133, 343
- Amphipoda 95, 681
- Amphiprion clarkii 629, 637–638
- Amphispiza bilineata 214
- Anabantidae 4, 10
- Anabas 11 *testudineus* 4
- Anablepidae 834
- Anacanthobatis 52, 54
- Anchovia 423, 427
- clupeoides* 427
 - macrolepidota* 427
 - surinamensis* 427
- Ancistrinae 1007
- Anguilla 221–223, 588
- Anguillidae 588
- Anguilloidea 73
- Anolis 315–323, 747, 831, 861
- aeneus* 755
 - aequatorialis* 320, 323
 - aquaticus* 320
 - attenuatus* 320
 - bourgeaei* 317–320
 - carolinensis* 316–318
 - carpenteri* 320
 - chloris* 320
 - chrysolepis* 714
 - cooki* 853–862
 - cristatellus* 318, 853–862
 - cupreus* 318–319
 - cupreus macrophallus* 320
 - cupreus spilomelas* 320
 - cuprinus* 316–320
 - dollfusianus* 320
 - dunni* 318, 321
 - evermanni* 860
 - gadovi* 318
 - gadovii* 321
 - garmani* 748
 - gemmosus* 318–319, 321
 - grahami* 748
 - gundlachi* 860
 - heteropholidotus* 319, 321
 - humilis* 317–318, 321
 - intermedius* 321
 - isthmicus* 316–319, 321
 - lemurinus* 319, 321
 - lemurinus bourgeaei* 316
 - “*lemurinus*” 321–322
 - limifrons* 321–322, 756, 826, 830–831
 - lineatus* 747–757
 - liogaster* 318
 - lionotus* 321
 - megapholidotus* 321
 - monensis* 860
 - nebuloides* 318–319, 321
 - nebulosus* 319, 321
 - nigrolineatus* 321–322
 - pentaprion* 321
 - peraceae* 321–322
 - polylepis* 321
 - princeps* 323
 - quercurum* 316–319, 321
 - sagrei* 826, 830
 - sericeus* 317–319, 321–322
 - sminthus* 321–322
 - subocularis* 318–319, 321
 - taylori* 318, 321
 - tropidolepis* 321–322
 - tropidonotus* 316–318, 321–322
 - valencienni* 748
 - villai* 317–318, 321, 323
- Antennaria 33
- Antennarii 28
- Antennariidae 27–44
- Antennarioidae 28
- Antennarioidei 27
- Antennarius 27–30, 33–34, 36–38, 41–42
- analis* 36
 - asper* 38
 - avalonis* 27, 36
 - bermudensis* 36
 - biocellatus* 28, 36
 - chironectes* 33
 - coccineus* 36
 - commersoni* 34, 39
 - cryptacanthus* 40
 - doreensis* 28, 36
 - drombus* 36
 - duescus* 34, 36
 - filamentosus* 70
 - hispidus* 34
 - horridus* 34
 - indicus* 28, 36
 - maculatus* 27, 34, 36
 - mitchelli* 39
 - multicellatus* 34
 - nummifer* 36
 - ocellatus* 36
 - pardalis* 36
 - paukiradiatus* 28, 36
 - pictus* 34–35
 - radius* 28, 36, 42
 - randalli* 36
 - rosaceus* 28, 36
 - sanguineus* 28, 36
 - sarasa* 36
 - senegalensis* 36
 - striatus* 27, 34
 - subteres* 36
 - tuberous* 28
 - zebrinus* 34
- Antennatus 27–30, 33, 36–38, 42
- bigibbus* 37
 - strigatus* 36–37
 - tuberous* 35, 37, 41

- Anthia sexguttata* 1016
Anthias 199 *squamipinnis* 200
Anura 302, 390, 868, 966
Aparallactus 644 *concolor* 642, 644
Apotopus 668–669 *lineatus* 669
Apereidon affinis 235
Aphyonidae 579
Aplocheilus 9–12 *dayi* 9
Apogon 80
 doderleini 189
 gracilis 189
Apogonidae 79, 189
Apterygocampus 567
Aranida 681
Ariosa 79
 anago 79
 mellissi 591
Arius 265–266 *gambiensis* 265
Arizona 216 *elegans* 216
Arrhyton 648
Artemia 88 *salina* 833
Artennarius 33
Arychnobatis 52
Ascaphidae 283
Ascaphus truei 283–284
Aspidites melanocephalus 959
Aspidorhynchus 500
Astyanax fasciatus 937
Aystasia gangetica 748
Atherina 9, 11–12, *presbyter* 9
Atherinomorpha 9–12, 16
Atherinomorus 11–12 (*Pranesus*) *endrachtensis* 9
Atherinops 9, 11–12
Antimora rostrata 540
Atractaspis 520, 645
Atractosteus 476, 486, 496
 “*Atractosteus*” 493
Atractus 646
Atriplex
 canescens 371
 polycarpa 371
Aulostomidae 1013
Australasomus 498
Aves 135
Avicennia nitida 854
- Bagarius* 265
Bagrus meridionalis 150
Balistidae 190, 235
Barathrites 540
Barathronus 579–584
 bicolor 579–584
 multidens 579–584
 pacificus 579
 parfaiti 579–580, 583
 sp. 579, 583
 unicolor 579–583
Barbus 265
Bassogigas 540
Bathymyrinae 72, 78–79
- Bathymyrus* 78–79 *simus* 78
Bathyraja 45–50, 52, 54
 asperula 45–58
 spinifera 45–58
 tobitukai 45, 49, 56
Bathysaurus ferox 540
Batrachopus 37, 668–669 *deweyi* 669
Belonesox 924, 928–929 *belizanus* 924, 929–930
Beloniidae 834
Beneckea harveyi 79
Benthocometes 540
Betula verrucosa 708
Bhanotia 567
Bivalva 681
Bivibranchia 4, 11–12
Blachea 79 *xenobranchialis* 73
Blattidae 219
Blenniidae 189
Blighia sapida 748
Bobasatrania 497
Bodianus
 axillaris 1016
 diana 1016
 mesothorax 1016
 neili 1016
Boraginaceae 814
Bouteloua sp. 815
Brachionichthyinae 28
Brachystis frenatus 997
Brachyphyllea carvernorum 219–220
Brachyrhaphis 924, 928–929, 930
Breviraja 52, 54 *stehmanni* 55
Bromus rubens 815
Brotula 1009
Brycon 1007 *guatemalensis* 937
Bryophyllum pinnatum 748
Brysonima coriacea 748
Bufo 18–19, 23, 1024
 americanus 18–26, 1023–1024
 americanus × *Bufo fowleri* 19, 23
 arenarum 23
 atrigularis 912
 boreas 23, 283–290, 673, 694–695, 701, 974
 boreas halophilus 689–696, 696–701
 bufo 1024
 calamita 1024
 canorus 974–975, 1023–1024
 cognatus 23, 1023–1024
 exsul 1024
 fowleri 18–26
 holdridgei 1024
 houstonensis 1024
 marinus 219
 punctatus 23
 quercicus 1024
 raddei 1024
 speciosus 23, 1024
 spinulosus 974
 stentor 912, 914
 terrestris 1023–1025
 typhonius 1024

- valliceps* 1023–1024
viridis 23
woodhousei 1023–1024
woodhousei fowleri 18
woodhousei australis 731
Bulbonaricus 565–571
brauni 565–571
brucei 565–571
davaoensis 565, 567–570
- Caiman**
crocodilus 843
sclerops 666
- Caligus** *hobsoni* 758
- Callichrous** *bimaculatus* 235
- Callinectes** *sapidus* 185
- Callionymus** 4, 11–12
- '**Callisaurus** *draconoides* 371, 826, 830
- Calloselasma** *rhodostoma* 770–772
- Cambarus** 864
- Camissonia** 815
- Campostoma** *anomalum* 134–135, 137, 446, 449, 451
- Candacia** 598
- Canella** *winteriana* 854
- Canobius** 497
- Capellaria** 37
- Carangidae** 190
- Caranx* sp. 198
- Carapidae** 410
- Carapus** 415, 419–420
cinereus 413, 415
owasianus 417, 419
rendahli 419–420
- Carassius** 265–266 *auratus* 133, 259, 992, 998
- Caretta** *caretta* 153, 471
- Carphophis** 641–645, 649, 777 *amoenus* 642–643
- Cassia** *emarginata* 748
- Catotomus** 3, 11–12, 1021 *plebius* 1021
- Centrarchidae** 149, 996
- Centriscidae** 1013
- Centropomidae** 144
- Centropomus** *ensiferus* 262
- Centropyge** 194
interruptus 197–198
potteri 197
spp. 197
- Cephalopholis**
pachycentron 190
urodelus 190
xanthopterus 189
- Cephaloxenus** 497
- Cetengraulis** 422–427
edentulus 422–426
juruensis 422–426
mysticetus 423, 425–426
- Chaetodon** 11–12 *miliaris* 7, 12
- Chalcinus** *elongatus* 263
- Chamaeleolis** 322
- Chaemaelinorops** 322
- Chaetodon** *auriga* 758–761
- Chanos** *chanos* 629
- Chara** 999–1000 *vulgaris* 151
- Characoidei** 13
- Chaunacinae** 28
- Chaunax** 28
- Chela** 11–12 *bacaila* 4
- Chelidae** 780
- Chelinus** *diagramma* 190
- Chelonia** *mydas* 153, 774–776
- Cheloniiidae** 546
- Chelydria** 465 *serpentina* 209, 546–550
- Chelydridae** 546
- Chironectiformes** 28
- Chironectes** 37–38
bifurcatus 68, 70
bougainvilli 40
filamentosus 40, 69–70
furcipilis 27, 37
trisignatus 38
variegatus 37
- Chironectus** 37
- Chironius** 642, 647
exoleatus 642, 646–647
fuscus 647
- Chironomidae** 218
- Chirotherium** 668–669 *lulli* 669
- Chlopsis** *bicolor* 589
- Chondrichthyes** 45, 495
- Chondropython** *viridis* 959
- Chrisostoma** 9, 11–12
- Chromis**
caerulea 189–190
punctipinnis 997
- Chrysemys** *picta* 546–547, 811–812, 818
- Chrysichthys** 264–266
filamentosus 265–266
nigrodigitatus 265–266
walkeri 265–266
- Chrysophyllum** *argenteum* 219
- Cichla** *ocellaris* 536
- Cichlasoma** 933
affine 937, 939
aureum 933, 936–939
bifasciatum 937
callolepis 933, 936–937, 939
champotonis 933, 937–939
citrinellum 534, 537
elliotti 936–939
helleri 933, 936–939
helleri meeki 939
hyorhynchum 937, 939
intermedium 937
maculipinnis 938
meeki 534–538, 933, 937–939
octofasciatum 937
pasionis 933, 938–939
salvini 937
severum 534
socolofi 933–940
(Thorichthys) 939

- (Thorichthys) socolofi* 939
urophthalmus 537
- Cichlidae, 6, 13, 149
Cirrhilabrus 200
Cirrhitichthys falco 190
Cirrhitidae 190
Cladocera, 85, 93, 95, 218, 681
Clarias 265–266
batrachus 265
lazera 265
Claudius 456–462, 466, 469–471, 474 *angustatus* 457–461
Clausocalanus spp. 598, 600
Cleidopus 79
Clelia 642, 644 *scytalina* 644
Clepticus parrae 195, 199
Clethrionomys gapperi 978–979
Clupea
harengus 357–364
pallasi 362
Clupeopetersius 82
Cnemidophorus 713–714, 718
lemniscatus 713–719
sexlineatus 826–832
sp. 214
tigris 371
uniparens 830
Coccocola uvifera 854
Coleonyx variegatus 763
Coleoptera 681, 755
Collembola 755
Colostethus nubicola 284, 288
Coluber jugularis 642, 648
Colubridae 215, 516, 520, 645
Colubrina arborescens 854
Conchocia spp. 598, 600–601
Congridae 72–73, 404
Congriniae 72, 79
Conia tenuis 642–643
Coniophanes 642–644
Conocarpus erectus 854
Conophis 641–646, 648–650 *vittatus* 642–643
Contia 641–643, 649
Copepoda 95, 681
Coregonus
artedii 879–880
clupeiformis 238–242
hoysi 878–883
reighardi 800
Coris variegata 190–191
Corixidae 218
Cortaderia attacamensis 727
Corycaeus 600
Corydoras 266
Coryphaenoides
carapinus 540
rupestris 540
Cosmocerca sp. 680
Cottus 949
asper 531
gulosus 949
klamathensis 949–959
klamathensis klamathensis 954–955, 957
klamathensis macrops 954–955, 957–958
klamathensis polyporus 951–952, 954, 956–958
macrops 949, 951
tenuis 949
Craterocephalus 9, 11–12
Crocodylus
acutus 153–161
niloticus 843
novaeguineae 154
Crotalus 213–215
adamanteus 978
atrox 213–215
horridus 976–981
mitchelli 213
mollassus 213–215
viridis 213, 976
Crotaphytus collaris 214
Croton californicus 371
Cruriraja 52
Cryophis 216
Cryptantha 813–814
Ctenogobiops pomastictus 189
Ctenolabrus 11–12
exoletus 7
rupestris 7
Ctenopharyngodon
idella 337–342
idella × *Hypophthalmichthys nobilis* 337–342
Ctenopoma 4, 11–12 *acutirostre* 4
Culicidae 218
Cuora 458, 463
Cyathopharynx furcifer 8
Cyematidae 588
Cymatogaster aggregata 223–225
Cymopterus 815
Cynops pyrrhogaster 739–742, 745
Cyprinidae 138, 149, 337, 385, 442–455, 617, 652, 990, 992, 966
Cyprinodon 58, 61–63
beltrani 61–64, 66–67
labiosus 61–63, 66
macularius 364–369, 823
maya 61, 66–67
nevadensis 291, 297
simus 61, 66
variegatus artifrons 63
verecundus 58–68
Cyprinodontidae 231, 364, 797, 834, 992
Cyprinoidei 3, 10–12, 16
Cyprinus 10–12, 14, 265–266
carpio 4–5, 142, 795, 800–803, 992
carpio × *Carassius carassius* 992
- Dactylobatis* 52
Dapedium 498
Dascyllus
albicella 758–759

- aruanus* 637
melanurus 637
reticulatus 190
scabra 642, 647
Decapterus sp. 198
Deirochelys 470 *reticularia* 812
Delphinium 815
Dendrobates auratus 284
Dendrophidion dendrophis 642, 647
Dermochelyidae 546
Dermochelys coriacea 552–553
Descurainia 815
Desmognathus 343–344, 346, 350, 353–355, 687
 aeneus 343
 auriculatus 343, 354
 brimleyorum 343
 fuscus 343–356, 684–688
 fuscus × *Desmognathus ochrophaeus* 344
 imitator 356
 monticola 343–344, 354, 684
 ochrophaeus 343–344, 353–354
 quadramaculatus 343–344, 354
 santeetlah 343
 weltneri 343
 wrightii 343
Diadophis 641–643, 649, 777 *punctatus* 642–643
Dibranchus 41
Dicrolene intronigra 538–541
Dionda 995 *episcopa* 995
Diplodactylus 1025–1028
 ciliaris 1026–1027
 elderi 1027
 michaelseni 1027
 rankini 1026–1027
 (*Strophurus*) 1025, 1027
 strophurus 1026–1027
 wilsoni 1027
Diplodus holbrookii 174–187
Dipnoi 495
Dipsadidae 645
Dipsadini 645
Dipsas 645
Dipsosaurus dorsalis 371, 830
Diptera 681, 755
Dipturus 52
Dispholidus typus 642, 647
Dromicus 645
Drosophila
 melanogaster 310
 mercatorum 718
Dryinoides 648–649
Drymarchon 776–777
 corais 642, 647
 corais couperi 770–778
Drymobius chloroticus 642, 647

Echinophryne 27, 29, 31, 38–39, 41–42
 crassispina 28, 35, 38–39
 glauerti 27, 38
Echiodon 410–422

anchipterus 410–422
cinereus 410–415, 419
coheni 410–422
crymargarites 410–413, 417, 421
dawsoni 410–413, 417, 420–421
dentatus 410–412, 414, 417, 420
drummondii 410–414, 417
exsiliens 410–413, 419, 421
owasianus 410–413, 417–419
rendahli 410–413, 419–420
 species 410–412, 421
Edalorhina 918 *pustulata* 911
Eigenmannia 136
Elapidae 520
Eleotridae 189, 237
Eleutherodactylus 766
 coqui 850
 cuaquero 766
 matudai 765
 melanostictus 765–768
 miles 765
 platyrhynchus 765
Elops 480
Emoia 571–578
 caeruleocauda 571–572, 574, 576
 cyanura 571–573, 577–578
 maculata 571–572, 574–575, 577–578
 maculata superspecies 571–578
 rufilabialis 571–578
 schmidti 571–575, 577–578
 sp. 575, 577
 spp. 572
 taumakoensis 571–578
Emydidae 546
Emydoidea 470 *blandingii* 782
Emys depressa 782
 “*Emys depressa*” 783
Enchelycore 542
 anatina 541–544
 lichenosa 542
Enchelyocampus 565–571
 brauni 565–566, 568–569
 sp. 566
Engraulis 422–427
 anchoita 422–423, 425–426
 australis 425–426
 (*Cetengraulis*) 425
 edentulus 422
 engrasicolus 425–426
 eurystole 423, 425–426
 guineensis 425–426
 japonicus 425–426
 juruensis 422–423
 mordax 422–423, 425–426
 mysticetus 422
 ringens 422–423, 425–426
Engystomops petersi 908
Ephedra 815
Ephemeroptera 681
Epibulus insidiator 1
Epirates

- angulifer* 219
inornatus 219-220
- Epidonidae** 1014
- Epinomen* 1014
- Epinephelidae** 189-190
- Epinophelus* 2, 793 *striatus* 793
- Equula elongata* 884-891
- Equus asinus* 702
- Eremophila alpestris* 214
- Eretmodus* 8, 11-12 *cyanostictus* 7-8
- Ericymba buccata* 386
- Erimyzon suetta* 150
- Erophyllo sezekorni* 219-220
- Erythrolamprus* 645
- Esocidae** 992
- Esox* 476, 481-482, 484, 491, 493-494 *lucius* 191, 205-208, 255-258
- Etheostoma**
- blennioides* 383
 - (*Catotomus*) 992
 - (*Catotomus*) sp. 992-994
 - kennicotti* 993
 - nianguae* 383
 - nigrum* 380, 383
 - radiosum* 378, 380-381, 383
 - spectabile* 291-301, 380, 383
 - squamiceps* 993
- Etomopterus spinax* 794-797
- Eucalanus* 598 spp. 598, 600, 605
- Euchaeta* 598 spp. 600-601
- Eucinostomus* 5, 11-12, 893-894, 896, 898-900
- argenteus* 893, 895, 897, 900-901
 - gula* 893, 895, 900-901
 - harengulus* 900-901
 - havana* 893
 - jonesi* 893, 895, 900-901
 - lefroyi* 893
 - melanopterus* 893
 - pseudogula* 895
- Eudontomyzon mariae* 503
- Eupemphis**
- freibergi* 908, 910
 - nana* 917, 919
 - paraensis* 908
 - ruthveni* 912-913
 - schereri* 903, 908, 911
 - trinitatis* 912
- Euphorbia punicea* 748
- Eupomotis** (*Lepomis*) 3
- Eurycea* 355, 428-434, 436-437, 439
- latitans* 428-440
 - longicauda* 744-745
 - lucifuga* 744, 863
 - lucifuga longicauda* 863
 - neotenes* 428-441
 - neotenes* × *Eurycea tridentifera* 430, 432-433, 437
 - sp. 429, 432-437
 - tridentifera* 428-441
 - troglodytes* 428-440
- Eutetideus* spp. 598
- Euthynnus*
- affinis* 1003-1004
- lineatus* 1000-1005
- Euxiphipops* 194
- Exocoetus monocirrus* 1001, 1003-1004
- Fagus silvatica* 708
- Farancia* 641-646, 649 *abacura* 642-644
- Felis domesticus* 219
- Ficimia publia* 642, 647
- Fierasfer* 420 *dentatus* 420
- Fistulariidae** 1013
- Forcipiger longirostris* 1
- Formicidae** 755
- Fowlerichthys* 33 *floridanus* 33-34
- Fundulus* 3, 7, 10-12, 59, 231-232, 820-821, 824
- bermudae* 820-821, 824
 - chrysotus* 9
 - confluens* 822
 - diaphanus* 133
 - grandis* 798-799, 820-821, 824-825, 836
 - heteroclitus* 117, 522-525, 798, 820-822, 824
 - heteroclitus heteroclitus* 524, 820-821, 824
 - heteroclitus heteroclitus macrolepidotus* 820-824
 - n. sp. 820-821, 824
 - olivaceus* 386
 - pulverus* 797-800
- Furo* 498
- Gadidae** 1008
- Gadus* 1009
- morhua morhua* 147
 - tenuis* 1008-1009
- Galaxea* 567
- clavus* 571
 - fascicularis* 569
 - musicalis* 569
- Galerinus orientalis* 1016
- Gambelia wislizenii* 371
- Gambusia* 530, 922, 924, 928-929, 931-932
- affinis* 108-119, 299, 386, 529-530, 924, 930, 989
 - amistadensis* 529-530
 - (*Arthrophallus*) 924
 - baracoana* 921, 931
 - bucheri* 921, 931
 - caymanensis* 921, 931
 - gaigei* 530
 - howelli* 911, 926, 931
 - hubbsi* 921, 931
 - lemairei* 924, 930
 - manni* 117, 921, 931
 - monticola* 921, 931
 - myersi* 930
 - nicaraguensis* 930
 - oligosticta* 921, 931
 - (*Orthophallus*) 924
 - puncticulata* 921-922, 926, 928-930
 - puncticulata baracoana* 922, 925-926, 928
 - puncticulata bucheri* 922, 925-926, 928
 - puncticulata manni* 922, 924-926, 928

- puncticulata monticola* 922, 925–926, 928
puncticulata puncticulata 921–922, 924–928, 931
puncticula yucatana 921–922, 924–927
sexradiata 58
vittata 924, 928–930
wrayi 930
yucatana 922, 928–931
- Gambusiini** 924, 929
- Garmanella* 235 *pulchra* 235
- Gasterosteoidae** 1013
- Gasterosteus* 6, 10–11, 454, 532
aculeatus 5–6, 87–97, 143, 258–262, 530–534, 795
dorissus 531–533
leiurus 87, 92
semiarmaatus 88
trachurus 87, 92
- Gazza* 884
achlamys 996
minuta 886
- Gekkonidae** 1025
- Genicanthus* 194–195, 199
bellus 199
caudovittatus 194, 199
lamarchi 194–200
melanospilos 195, 197
semifasciatus 195, 197, 199
spp. 194–195, 197, 199
watanabei 199
- Geochelone* 702, 704–706
elephantopus 704
elephantopus ephippium 704–706
elephantopus porteri 704–706
elephantopus vandenburghi 702–707
gigantea 702, 704–705
- Geophis* 646
- Geotria australis* 503, 511–513
- Gerreidae** 5, 11–12, 893
- Geukensia demissa* 522–524, 821
- Gila**
cypha 790–793
elegans 790–793
- Ginglymodi** 496
- Ginglymostoma* 203
- Glaucomys volans* 978
- Glossogobius* 11–12 *giuris* 4
- Glypidodontops**
flavipinnis 190
rollandi 189
- Gnatholepis* 227–228 *hendersoni* 227–229
- Gnathophis* 79
- Gobiidae** 189–190, 235, 237
Gobio 4, 10–11, 265 *gobio* 4
- Gobiodon* 237
- Gobionellus shufeldti* 235–238
- Golem* 40
- Goodeidae** 102, 834, 836
- Gopherus polyphemus* 818
- Gorgasia* 404, 406, 408
barnesi 404–409
hawaiiensis 404–406
japonica 404–406
- maculata* 404–406
naeocepaeus 404–406
preclara 404–406
punctata 404–406, 409
sillneri 404–406
- Grammatidae** 1016
sexlineatus 1015–1017
sexlineatus proerytheraeus 1015
- Grammatidae** 1015–1016
- Graptemys flavimaculata* 818
- Grayia* 815
- Guaiacum officinalis* 748
- Gurgessiella* 45, 47–50, 52–56
atlantica 57
atripinna 57
cubensis 57
dorsalis 57
(Fenestraja) 43, 52–53, 55–57
furvescens 55–57
(Gurgessiella) 45, 52–56
ishiyamai 57
sibogae 57
sinusmexicanus 57
- Gymnocephalus* 11–12 *cernua*, 7, 14
- Gymnotorax**
funebris 589
javanicus 251
moringa 589
- Gymnotiformes** 134
- Gyrinophilus* 355
- Haemulidae** 190
- Halecomorphi** 496
- Halecostomi** 496, 499
- Halichoeres hooveri* 189–192
- Haplochromis*
burtoni 14
pictus 150
pleurostigmoides 150
- Haplolepididae** 497
- Helostoma* 11–12 *temmincki* 4
- Hemiptera** 681
- Hemirhamphidae** 834
- Hemitremia flammæ* 901
- Heterandria* 720–721, 725
formosa 720–726
sp. 937
- Heterenchelyidae** 588
- Heterocongrinae** 404
- Heterodon* 641–647, 649 *platyrhinos* 642
- Heterotilapia multispinosa* 534
- Hilaria rigida* 815
- Hildebrandichthys* 422 *setiger* 422–427
- Hippolytidae* sp. 80
- Hippolytidæ* 80
- Histionotus* 498
- Histiophryne* 21, 28–30, 40–42
bougainvilli 28, 40
cryptacanthus 35, 40
scorteia 27, 39

- Histro* 27–30, 32–34, 37, 41–42 *histrio* 28, 37
- Holacanthus*
 passer 195, 199
 tricolor 197
- Hollardiinae* 6–7, 11–12
- Homoptera* 755
- Hoplochelys* 469–470
- Horaichthys setnai* 833–839
- Hyalella azteca* 999
- Hybognathus* 453
 hayi 446, 449, 452–453
 nuchalis 446, 449, 453
 placitus 446, 449, 451–453
- Hybopsis*
 aestivalis 446, 449, 452–453
 amblops 446, 449, 453
 x-punctata 446, 449, 453
- Hydrilla* 721
- Hydrurga leptonyx* 787
- Hyla* 609–616
 arborea 609
 arborea arborea 609–615
 arborea sarda 609–615
 chinensis 609–615
 chrysoscelis 844–852
 cinerata 731
 crucifera 302–308
 ebraccata 731
 hallowni 609
 japonica 609–615
 meridionalis 609–615
 regilla 727–732
 savignyi 609–615
 suweonensis 609–615
 versicolor 733–738
- Hylidae* 302, 390, 609, 613, 868
- Hynobius* 866
- Hyperentelium nigricans* 386
- Hyperolidae* 283
- Hypomesus pretiosus* 1004
- Hypophthalmichthys* 340
 molitrix 340
 nobilis 337–342
- Hypostominae* 1007
- Hypsilema ochrorhyncha* 763
- Hypsipops rubicundus* 761
- Hystericarpus traski* 223–227
- Ichthyocampus* 565–566
 (*Bulbonaricus*) 565
 (*Bulbonaricus*) *davaoensis* 565–567
 davaoensis 565–566
 fossor 513
 gagei 502–515
 greeleyi 511–512
 hubbsi 511
- Ictaluridae* 873
- Ictalurus* 97–101, 265–266, 439, 873–878
 (*Amiurus*) *melas* 100
 catus 873, 875–877
- catus* × *Ictalurus furcatus* 873, 875–876
 furcatus 873–877
 furcatus × *Ictalurus punctatus* 873–876
 hazenensis 531
 (*Ictalurus*) *lupus* 100
 (*Ictalurus*) *punctatus* 100
 lupus 97–101
 melas 97–101
 punctatus 97–101, 550, 873–877
 punctatus × *Ictalurus catus* 873, 875–876
 punctatus × *Ictalurus furcatus* 873–877
- Idus* 4, 11–12, 14 *idus* 14
- Iguana iguana* 826
- Ilyodon* 102–107
 furcidens 102
 xantusi 102
- Imantodes* 215
- Isoetes* 673
- Isopoda* 95
- Itigibius ornatus* 189
- Ixora*
 coccinea 748
 parviflora 748
- Juniperus communis* 708
- Justicia* sp. 379
- Kaloula pulchra* 283
- Kanazawaichthys* 34 *scutatus* 34
- Kassina senegalensis* 283
- Katsuwonus pelamis* 1004
- Kaupichthys hyoprorooides* 589
- Kinosternidae* 456, 458–459, 469, 471–472, 474, 546
- Kinosterninae* 456, 461–462, 469, 471, 474
- Kinosternon* 456–460, 462–468, 471–474
 alamosae 456
 angustipons 456–457, 463, 472–473
 bauri 456–457, 463, 466, 473
 dunni 456, 463, 472
 flavescens 456–457, 464, 466–467, 473
 flavescens spooneri 818
 herrerae 456–457, 463–464, 472–474
 hirtipes 456–457, 464, 473
 integrum 456–457
 leucostomum 456–457, 463–465, 468–469, 473
 oxaceae 456
 scorpioides 456–457, 463–469, 472–473
 sonoriense 456–457
 subrubrum 456–457, 573
- Krameria* 815
- Kuiterichthys* 27–29, 32, 37–38, 41–42 *furcipilis*
 28, 38
- Labridae* 189–190, 245, 758
- Labrodes phthisiophagus* 758–762
- Labrus* 11–12
 bergylta 7
 bimaculatus 7
- Lacertilia* 826

- Lagodon* 11–12 *rhomboides* 7, 174–187
Laguncularia racemosa 262, 854
Lampetra
 appendix 513
 fluvialis 503, 511–513
 japonica kessleri 503, 511
 planeri 503, 511–513
 richardsoni 503, 511
 tridentata 511, 513
Lampropeltis 216 *pyromelana* 216
Lamrophis 644 *fuliginosus* 642, 644
Lantana 963
Larrea 813, 815 *tridentata* 371, 812
Larus *fuliginosus* 786
Lates 5, 11–12
 calcarifer 144–149
 niloticus 5
Lebistes *reticulatus* 140–141
Leiognathidae 81, 884, 891
Leiognathus 884
 aureus 886
 berbis 886, 891
 binuds 886
 brevirostris 886
 daura 886
 edwardsi 886, 891
 elongatus 884–892
 equulus 886
 fasciatus 886
 leuciscus 886, 891
 lineolatus 891
 moretoniensis 886, 888, 890–891
 nuchalis 886
 popei 880, 884, 888
 rapsoni 888
 rivulatus 886
 smithursti 886
 splendens 886
 stercocarius 884–891
Lepidocephalichthys 265
Lepidorhynchus 79
Lepidotes 498–499
Lepisosteus 476–502, 649
 (*Astropterus*) 476
 (*Lepisosteus*) 476
 oculatus 477, 493
 osseus 149–152, 476–483, 485–490, 492, 494
 platostomus 476–479, 481–489
 spatula 477–478, 481, 485–489, 492
Lepomis 11–12, 14, 439
 cyanellus 149, 385, 439
 gibbosus 260
 macrochirus 162, 386, 996–999
 megalotis 386, 997
 spp. 439, 550
Leptodactylidae 765
Leptodactylinae 919
Leptodactylus 765
 bufonius 780
 fuscus 778–780
 pentadactylus 778
Leptodeira 642–645 *septentrionalis* 644
Leptothrix *ahaetulla* 642, 647
Leucoraja 52
Liasis *fusca* 959
Limacina spp. 599–600
Limpus 552–553
Liophis 645
Litoria *ewigi* 868–873
Lobochilotes *labiatus* 14
Loigia *opalescens* 1001, 1003–1004
Lophiiformes 27
Lophiocharon 27–30, 32, 38, 41–42
 broomensis 38
 caudimaculatus 39
 lithinostomus 29, 39, 41
 trisignatus 28, 39
Lophius 37
 chironectes 33
 commersonianus 33
 histrion 37
 laevigatus 37
 pictus 33–34
 striatus 33
Loricariidae 1005, 1007
Lotus 815
Lubbockia 600
Lucania *parva* 230
Luciocephalus 11–12 *pulcher* 6–7, 13
Lumiconger 72–81 *rafaura* 72–81
Lupinus 815
Lycenchelys
 paxillus 540
 verrilli 540
Lycium 815 *andersonii* 812
Lycodontis *unicolor* 543
Lystrophis 648

Macilentichthys
 elongatus 885
 popei 885
 stercorarius 885
Macroclemyx 471
Macrodonogobius 227–229 *wilburi* 227–229
Macropodus 4, 11–12 *opercularis* 4
Macrorhamphosa 1013
Macrorhamphosidae 1013
Macrorhamphosus 595–596, 605–607
 elevatus 595, 603, 605
 gallinago 595
 gracilis 595–596, 598, 601–602, 606
 japonicus 596
 lancifer 595, 603, 605
 robustus 595
 sagifue 595
 scolopax 595–608
 scolopax var. *elevatus* 595
 velitaris 595
Macrosemiidae 484
Macrosemius 498 *rostratus* 484
Macrouridae 79

- Malacoraja* 52-55
fuliginea 55
kreffti 55
nigerrima 55
purpureiventralis 55
senta 55
spinacidermis 55
Malacothrix 815
Malapterurus 265-266 *electricus* 265
Mammalia 133
Mangifera indica 748
Marconus 266
Masticophis taeniatus 772
Mastigodryas melanolumus 642, 647
Mauremys
caspica rivulata 216-219
leprosa 217
Megupsilon 235 *aporus* 235, 238
Mehelya 644 *crossi* 642, 644
Meiacanthus atrodorsalis 189
Melichthys bunivira 591
Menidia 5, 11-12 *beryllina* 544-545
Mentzelia 815
Micralestes 263
Microhyliidae 283
Micropterus 439
dolomieu 149-152
salmoides 13, 117, 149, 550, 997
spp. 439
Microsetella spp. 600
Microtus pennsylvanicus 979
Monacanthus chinensis 191
Monocentridae 79
Monocirrhus 11-12 *polyacanthus* 4-5
Monomittopis 540 *agassizi* 540
Monophyllus redmani 219-220
Mordacia praecox 511
Morelia 960 *spilote* *variegata* 960
Moringuidae 588
Mormoops blainvillii 219
Morone 793 *saxatilis* 793-794
Mugil 3, 11-12 *cephalus* 996-997
Mullidae 189-190
Muraena
ansatina 542
helena 543
santaeheleneae 543
unicolor 591
Muraenidae 585, 585-593
Muscelus 203 *canis* 803
Myliobatis 203
Myoxocephalus
quadricornis 264
scorpius 264
Myroconger 585, 587-593 *compressus* 585-594
Myrocongridae 585, 589-593
Mystus 265
Nama demissum 815
Nandidae 4-5, 11-12
Nandus 11-12 *nandus* 4
Nannocalanus 598, 600 spp. 598
Nannocampus 567
Nanochromis 82-83, 86
consortus 83, 85
dimidiatus 82-83
minor 83, 85
nudiceps 82, 85
parilus 83-86
sp. undet. 83
splendens 83-86
squamiceps 83
transvestitus 82-86
Natrix
natrix 707-713, 826
tessellata 516-521
Nemachilus 265
Nematoscelis microps 250
Nemichthysiidae 588
Nemipteridae 190
Neobythitinae 538
Neochinorhynchus sp. 680
Neognathi 12
Neopomacentrus azyron 189
Neopterygi 500
Neoraja 45, 47-50, 52-53, 55
caerulea 56
(Fenestraria) 45, 48-50, 52, 54-55
(Neoraja) 45, 49-50, 54-55
stehmanni 56
Nerium oleander 748
Nerodia 644, 777
fasciata 642, 644
rhombifera 550-552
sipedon *sipedon* 865
Ninia 646
Nocomis 652-653, 656
asper 446, 449
leptocephalus 446, 449
micropogon 652-658, 990-992
micropogon \times *Rhinichthys cataractae* 652, 657, 990-991
platyrhynchus 653
Notagogus 498
Notimigonus crysoleucus 134-138, 149, 233, 446, 449,
452-453
Nothopsini 645
Notiocampus 567
Notophthalmus viridescens 162, 739-745
Notropis 137, 232, 385, 388-389, 451-452, 995
analostanus 995
atherinoides 446, 449
bairdi 446, 449
bifrenatus 995
boops 443, 446, 449
buchananii 446, 449
camurus 386, 446, 449
chalybaeus 995
chryscephalus 386, 446, 449, 617-628
cornutus 150, 617-628
cornutus chryscephalus 617

- emiliae* 134-135, 137, 446, 449, 991
fumeus 446, 449
galacturus 995
girardi 446, 449
greenei 446, 449
hudsonius 205-206
longirostris 386, 446, 449, 453
lutrensis 134-138, 233-234, 386, 447, 449, 453
(Luxilus) 617
(Lythrus) 385
maculatus 995
mekistocholas 995
nubilus 443, 447, 449, 994-996
(Opsopoeodus) emiliae 134
ortenburgeri 447, 449
perpallidus 447, 449
pilsbryi 447, 449, 994
potteri 447, 449
roseipinnis 386, 447, 449, 453
rubellus 447, 449, 995
sabinae 230
shumardi 447, 449
stramineus 447, 449
texanus 447, 449, 452
umbratilis 149, 385-390, 447, 449
umbratilis cyanoccephalus 385
umbratilis umbratilis 385
venustus 134-138, 233, 386, 447, 449
volucellus 447, 449
whipplei 447, 449, 995
zonatus 994
Nudiantennarius 27-29, 33, 36, 41-43 *subteres* 28, 36
- Odonata** 681
Odontobutis obscura 237
Okamejei 52
Oligodon sp. 642, 647
Oncaea spp. 598, 600
Oncorhynchus 485
kisutch 205-206, 208, 882
masou 148
tshawytscha 882
Ondatra zibethicus 550
Onuxodon 413
Ophidium 420 *dentatum* 420
Ophiinae 645
Opsopoeodus 137
Opuntia 814
echinocarpa 812
ramosissima 812
Orconectes 355, 864
Oryzias latipes 836
Osmerus mordax 882
Ostariophysi 3, 12
Ostracoda 95, 218
Otolithus (Gadus) tenuis 1009
Otozoum 669
Oxybelis fulgidus 642, 647
Oxyjulis californica 758, 761, 997
- Oxyporhamphus micropterus* 1001, 1003-1004
Oxyrhopus 645
- Paleoheterodon* 648-649
Paludicola pustulosa 912
Paralichthys lethostigma 981-988
Parapercidae 190
Parapercis cylindrica 190
Parapriacanthus 80
Parexocoetus 5, 11-12
Parodontidae 235
Parupeneus
cyclostomus 190
trifasciatus 189-190
Pavoraja 45, 47-50, 52-55
alleni 56
asperula 56
nitida 56
(Pavoraja) 45, 53, 55-56
spinifera 56
Pavorajini 45, 52, 55
Pectocarya 813-814
Pegasidae 1011-1012
Pegasoidea 1012-1013
Pegasus papilio 1012
Pelamis 786 *platurus* 786-789
Pelobatidae 289
Peltopleuridae 497
Pelusios 464
Penetopteryx 567
Penopus 540
Perca 5-6, 11
flavescens 151, 205-207
fluviatilis 5-6, 10, 147, 191
Percichthyidae 793
Percidae 291
Perciformes 630, 884, 1009
Percina
caprodes 380
copelandi 380, 383
nigrofasciata 299
pantherina 378-384
phoxocephala 383
sciera 380
Percomorpha 4, 9-10, 12, 16
Perleididae 497
Peromyscus
leucopus 978
sp. 979
Petromyzon marinus 503, 511-513, 879
Petrotilapia 11-12, 14 *tridentiger* 8, 14
Phallostethidae 1014
Phanerodon
atripes 997
furcatus 997
Phenacobius mirabilis 447, 449, 453
Philodryas 642-645 *chamissonis* 644
Philothamnus angolensis 642, 647
Pholidophoropsis 498
Pholidophorus 498-499
Pholidopleuridae 497

- Phoxinus* 991
eos 992
eos × *Phoxinus neogaeus* 992
erythrogaster 447, 449, 452
neogaeus 992
phoxinus 15, 998
Phractolaemus 4
Phrynelox 33
Phrynops
geoffroanus 783
tuberculatus 209
Phrynosoma m'callii 371
Phrynus palmatus 219
Phycis 1009
blennoides 1009
chesteri 540
simplex 1009
Phyllidia sp. 1016
Phylloctetus lanei 763
Phyllophryne 27, 31, 39–42 *scorteia* 28, 40
Phymatophryne 34
Physalaemus 902, 904, 907, 910–911, 914, 917–919
albonotatus 903, 914, 916–917
biligonigerus 903, 911, 914–917, 919
centralis 903, 914, 916
coloradorum 902–909, 911–912, 914–920
cuvieri 903, 911, 914, 916–917, 919
enesefae 903, 912, 914, 916, 918–919
ephippifer 903, 910, 914–916, 918–919
fernandezae 916, 919
freibergi 902–903, 911, 915
fuscomaculatus 917
gracilis 903, 914–916
henseli 916, 919
maculiventris 903, 914–916
nanus 903, 914–917, 919
nattereri 903, 911, 914–917, 919
obtectus 915
olfersi 916
paraensis 902, 915
petersi 902–906, 908–912, 914–919
pustulatus 902–906, 908–909, 911–912, 914–916,
 918–919
 “*pustulatus*” 907, 912, 917
pustulosus 780, 902–906, 908–919
riograndensis 916
santafecinus 916
schereri 902, 911, 915
signiferus 903, 914–917
stentor 902, 910, 915
Picea abies 708
Pimelodus lupus 97
Pimephales
notatus 386, 447, 449
promelas 447, 449, 453, 550
tenellus 447, 449
vigilax 134–136, 138, 233, 386, 447, 449, 452
Pisces 72, 133, 235, 291, 337, 385, 410, 442, 528,
 565, 579, 585, 652, 720, 893, 921, 990, 1008,
 1015
- Pituophis melanoleucus* 771
Platemys
macrocephala 782
pallidipictoris 782
radiolata 780–786
radiolata spixii 783
spixii 780–786
 “*spixii*” 782
Platuronides 251
Platyagium 497
Platysternon 471
Plectrohyla 390–397
avia 395
dasyuspis 390–391
glandulosa 390–391, 393, 395
guatemalensis 390, 394–395, 397
hartwegi 390, 394–397
lacertosa 390–391
pokomchi 390–397
pychnochila 390–391, 393
quechii 396
tecunumanii 390–397
Plectropomus leopardus 190
Plectrohynchus chaetodonoides 1016
Plethodon 355, 686
cineraceus 309–314, 686–687
hoffmani 687
jordani 739–740, 742–745
punctatus 687
Plethodontidae 309, 343, 739–747
Pleurodeles waltli 739–743, 745
Pleurodema 916, 918
Pleurodira 780
Pleuromamma 599–600
abdominalis 598, 600
spp. 599
Pleuronectes 11–12 *platessa* 5–6
Plumantennarius 38
Poecilia
latipinna 252–255, 836, 838
mexicana 937
picta 262
reticulata 140–144, 795
Poeciliidae 169, 252, 720, 836, 921
Poeciliinae 928–929
Poeciliopsis 169–174, 989
lucida 169–170, 173
monacha 169
Polychrus 322
Polydactylus virginicus 262
Polyodon 201, 649
feuillei 201
folium 201
spathula 201–204
Polypterus 476, 482, 484–485, 493, 496–497
Pomacanthidae 190
Pomacanthus 194 *imperator* 197
Pomacentridae 189–190, 630, 637–638
Pomacentrus
amboinensis 189–190, 192

- bankanensis* 189
coelestis 189–190
flavicauda 637
lepidogenys 189
popei 189–190
 sp. 190
wardi 637
Pontoporia 881
Porichthys 80
Porogadus 540
Powelliichthys ventriosus 592
Priapella compressa 937
Procambarus 385
Propterus 498
Prosopis glandulosa 371
Prunus spinosa 711
Psammobatis 52, 54
Pseudechis porphyriacus 772
Pseudemys
 — *floridana* 209
 — *scripta* 209–210, 812
 — *scripta scripta* 210
Pseudoboini 645
Pseudochromidae 189
Pseudochromis mccullochi 189
Pseudoeryx 644
Pseudopaludicola 918–919 *falcipes* 919
Pseudophryne 24
Pseudoraja 52, 54
Pseudotriton 355 *ruber* 866
Pseustes 645 *poeilonotus* 642, 647
Pteraichthys 668
Ptereotris evides 189
Pterolebias 528
 — *hoignei* 528
 — *longipinnis* 528–529
Pteronisculus 497
Pteronotus fuliginosus 219
Pterophryne 37
Pterophrynoidea 37
Pterophrynum 37
Pterophyllum 6, 11–12
Pterygoplichthys 1005–1008 sp. 1006–1007
Ptychocheilus lucius 790–793
Ptycholepis 497
Pygoplites 194
Pylodictus olivaris 997
Pyramodon 413
Pyramodontinae 413
Python 960, 964
 — *amethystinus* 959
 — *curtis* 959
 — *molurus* 959, 964
 — *molurus bivittatus* 963
 — *reticulatus* 964
 — *sebae* 964
 — *spilotus* 959
 — *spilotus spilotus* 959–965
 — *spilotus variegata* 960, 963–964
 — *timorensis* 959
- Quercus*
 — *prinns* 976
 — *robur* 708
 — *rubra* 976
- Raja* 46, 52
 — (*Atlantoraja*) 55
 — (*Atlantoraja*) *castelnau* 51
 — (*Atlantoraja*) *cyclophora* 51
 — (*Atlantoraja*) *platana* 51
 — (*Malacoraja*) 45, 47–50, 54–55
 — *nitida* 56
 — *plutonia* 57
 — (*Rioraja*) 55
 — (*Rioraja*) *agassizi* 51
 — *spinacidermis* 55
 — *waitei* 51, 55
- Rajella* 52
Rajidae 45, 52, 55
Rajoidei 45, 52, 55
Ramphosidae 1012
Rana 284, 398, 1020
 — *aurora* 1018
 — *aurora aurora* 1018–1022
 — *aurora draytoni* 1018–1022
 — *boylei* 1021
 — *boylii* 398
 — *calamitans* 211
 — *cascadae* 673, 1021
catesbeiana 162, 212, 283, 694, 727, 730, 974
chiricahuensis 940–943, 945, 947
clamata 210
clamitanus 162, 210–213, 289, 731
draytoni 1018
fisheri 940, 943, 947–948
forreri 945–947
halecina 398
johni 398–403
maculata 398
magnaocularis 940–947
moorei 398
muscosa 966–976, 1021
nigromaculata 609
onca 940, 943, 947–948
palmipes 398, 400–401
palustris 162
pipiens 289, 398, 869, 871–872, 940, 942–943, 945–947, 974–975
pipiens fisheri 943
pretiosa 1021
pueblae 398–402
pustulosa 398–402
septentrionalis 212
sierramadrensis 400
sinaloae 398–399
sphenocephala 289, 550
sylvatica 162, 166, 283, 871–872
tarahumarae 398–403
trilobata 398

- utricularia* 289
yavapaiensis 940–948
zweifeli 398–403
Ranidae 966
Ranodon sibiricus 866
Redfieldiidae 497
Regina 777
Reptilia 133, 660, 1025
Rhabdosargus holubi 185
Rhadinea 777
Rhadinophanes monticola 215
Rhamdia guatemalensis 937
Rhamphiphis 644 *oxyrhynchus* 642, 644
Rhinichthys 652, 658
 bowersi 652, 990–991
 “*bowersi*” 652–659, 990–992
 catarractae 652–658, 990–992
Rhinoclemmys aerolata 209
Rhizophora mangle 262
Rhycherus 27, 29–31, 40–42, 68–72
 bifurcatus 68
 filamentosus 28, 40, 68–71
 gloveri 69–72
 wildii 40, 68–70
Rhynchoconger 72, 79
Rhynchosauroides 669
Rivulichthys luelingi 528–529
Rivulidae 528
Rostroraja 52
Rubia tinctoria 989
Rubus fruticosus 708, 711
Rutilus rutilus 147
- Saccarius* 33 *lineatus* 33
Saccopharyngoidea 73
Salamandridae 739–747
Salix sp. 967
Salmo 480, 482, 484–485, 491, 493–494, 673
 aguabonita 120
 clarki 531, 672
 clarki seleniris 128
 gairdneri 120, 124, 129, 131, 264, 672
 newberryi 120, 129–130
 newberryi kamloops 129
 salar 148
 sp. 531
Salmonidae 992
Salvelinus 673
 fontinalis 147, 672
 lucoamaenis 148
 namaycush 882
Sarda chiliensis 1004
Sardinia pilchardis 297
Sargassum 42
Sarotherodon
 galilaeus 134
 mossambicus 147
Sauria 713
Saurichthyidae 497
Saurida gracilis 190–191
Scaphiodontophis zeteki 642, 647
Scaphiopus 778 *couchii* 289
Scaptochelys agassizi 811–820
Scaridae 190
Schismus barbatus 815
Sciaena 11 *cotter* 4, 12
Scincella lateralis 763
Scirpus validus 256
Scomber
 japonicus 1004
 scombrus 299
Scomberoides lyasan 190
Sebastes 3, 11–12 *marinus* 5
Secutor 884
 indicus 886
 insidiator 886
 ruconius 886
Seミonotus 499
Semitilapia atromaculatus 447, 449, 453
Sergestes sp. 250
Serpentes 660
Serranidae 793
Serrivomer spp. 251
Serrivomeridae 249
Siluridae 235
Siluroidei 1005
Simophis 645 *rhinostoma* 642, 647
Siokunichthys 565 (*Bulbonaricus*) 565
Siphonia 79
Snyderidria 413
Solenostomidae 1012
Spartina 798
Spathodus 8, 11–12 *erythrodon* 7
Spectrunculus 540
Sphaerodactylus
 levinsi 769–770
 mariguanae 769
 micropithecus 768–770
 monensis 768–770
 savagei 769
Sphaerooides 11–12 *oblongus* 7
Sphyraena viridensis 541
Sphyraenops 1014
Spilotes pullatus 642, 647
Spongilla 82, 85
Spratelloides sp. 190
Squalus
 spathula 201
 Spatula 203
Staurotypinae 456–457, 459, 474
Staurotypus 456–462, 465–466, 469–471, 474
 salvini 457
 triporcius 457, 461, 471
Stegastes fasciolatus 629–640
Stegmosuchus 668
Stemonidium hypomelas 249–252
Stenorhina frenimvillei 642, 647
Stephanolepis cirrifer 235
Sternotherus 456–466, 469, 471–474
 depressus 818

- minor* 457, 462, 465, 472–473
odoratus 457, 462, 471, 546
Stethojulis bandanensis 189
Stipa speciosa 815
Stizostedion vitreum 191
Stylocheiron
 abbreviatum 250
 carinatum 250
 longicorne 250
 spp. 250
Suriana maritima 854
Swietenia mahagoni 748
Sylvilagus floridanus 978
Sympodus 11–12 *melops* 7
Syngnathia 52
Synaptilichnum 668–669 *pseudosuchoides* 669
Syngnathus 1012–1014
Syngnathidae 565, 1012
Syngnathoidei 1011, 1013–1014
Synodontidae 191
Synodontis 265–266 *schallii* 265
Synodus
 englemani 187–194
 intermedius 191
Syringodium filiforme 176–177, 183–184
Szygium jambos 748
- Tamarix aphylla* 371
Tamias striatus 978
Tantalophis 215–216, 645 *discolor* 215–216
Tantilla 645 *melenocephala* 642, 647
Tantillita 645
Taricha
 granulosa 672–683
 torosa 744, 863
Tarpon atlanticus 262
Tathicarpus 27, 29–30, 40–43 *butleri* 28, 49–51
Tautoga onitis 297
Tautogolabrus adspersus 297
Tectibranchiata 1016
Teiidae 713, 826
Teleogramma gracile 84
Teleostei 102, 245, 496
Telescopus semiannulatus 642, 647
Terrapene 463 *carolina* 818
Tetrahymena 744
Tetraodontiformes 7
Texasophis 660–661
 fossilis 660–661
 wilsoni 660–661
Thalassia testudinum 176, 184
Thalassoma
 amblycephalus 189
 duperreyi 245–248, 758–759
 lucasanum 200
 lunare 189
Thamnophis 516, 777
 sirtalis 865
 sp. 979
Thelotornis capensis 642, 647
- Thespesia populnea* 748
Thorichthys 938
 elliotti 938
 helleri 938
Thrasops jacksoni 642, 647
Thryssa rastrota 423
Thysanopoda
 aequalis 250
 monocantha 250
 spp. 250
Tilapia mariae 534
Tinca 265
Triacanthodinae 6, 11–12
Triantennatus 34
Trichechus manatus 996
Trichophryne 27, 29, 31, 38–39, 41–42 *mitcheli* 28, 35, 39
Trichoptera 681
Trimorphodon 642, 645, 647 *biscutatus* 642, 646–647
Trionyx 472, 649
Triportheus 263 *elongatus* 262–263
Triturus vulgaris 739–743, 745, 866
Tropidoclonium 777
Tropidodiplos 646
Tubificidae 218
Typhlomolge rathbuni 432
- Ulaema* 893
Uma 370–377 *inornata* 370–378
Umbra limi 133
Uniantennatus 34
Uranidea 949
 minuta 949
 tenuis 949
Uranoscopidae 1009
Uranoscopus kaianus 1009–1011
Urophycis 1009 *musicki* 1009
Urosaurus graciosus 371
Uta stansburiana 371, 812, 817
- Varanus komodoensis* 662–671
Vipera
 berus 771–772
 palaestinae 516, 520
- Walterinnesia* 520
- Xenochelys* 460, 469–471, 474
 formosa 470–471
 n. sp. 457
Xenocongridae 585, 588–593
Xenodon 643
Xenodontidae 645
Xenodontinae 645
Xenodontini 645
Xenophrynichthys 40
Xenopus laevis 844

Xiphophorus 9, 11-12, 924
 helleri 9, 937
 maculatus 148, 252
 variatus 252
Xyrauchen texanus 790-793

Xyelacyba 540
Zalophus californianus wollebecki 787
Zebrasoma flavescens 758-761
Zostera marina 821, 823

SUBJECT INDEX

COPEIA, 1984, Nos. 1-4

ACTIVITY, *Bufo boreas* (nocturnal activity & ambient light) 283-290; *Desmognathus fuscus* (nocturnal activity & moisture) 684-688; *Bufo boreas halophilus* (activity patterns & thermal biology) 689-696; (temperature & daily emergence time) 696-701; *Natrix natrix* (movement, home range, habitat use) 707-713.

AUTOTOMY, *Phyllodactylus lanei* (tail autotomy inhibits tonic immobility) 763-764.

BEHAVIOR, *Micropterus dolomieu* (brood care of *Lepisosteus osseus*) 149-152; *Synodus englemani* (predatory behav.) 187-194; *Genicanthus lamarchi* (reprod. behav., social organization) 194-200; *Bufo boreas* (nocturnal activity) 283-290; *Plethodon cinereus* (agonistic behav.) 309-314; kinosternid turtles (behav. in relation to shell kinesis) 356-375; *Fundulus heteroclitus* (spawning site selection) 522-525; *Cichlasoma meeki* (color pattern variability & behav.) 534-538; *Hyla regilla* (male mating behav.) 727-732; *Labrodes phthirophagus* (preference for ectoparasite-infected host) 758-762; *Phyllodactylus lanei* (tail autotomy inhibits tonic immobility) 763-764; *Calloselasma rhodostoma* (combat ritual) 770-772; *Agiistrodon piscivorus* (sexual defense) 772-774; *Leptodactylus fuscus* (foam production) 778-780; *Fundulus (bermudae, grandis, h. heteroclitus, heteroclitus macrolepidotus, n. sp.)* (reprod. behav. of spp. from east coast US & Bermuda) 820-825; *Anolis (cooki, cristatellus)* (competitive interference) 853-862; *Rana muscosa* (behav. thermoregulation) 966-976; *Crotalus horridus* (foraging behav.) 976-981; *Natrix nivalis* (spawning activity) 994-996; *Lepomis macrochirus* (cleaning behav.) 996-998; (group size & feeding rates) 998-1000; *Rana (a. aurora, aurora draytoni)* (behav. differences) 1018-1022; *Bufo terrestris* (amplexus displacement) 1023-1025.

BLOOD, sampling of, from turtles 209-210; *Gila (cypha, elegans)*, *Ptychocheilus lucius*, *Xyrauchen texanus* (hematology of) 790-793.

BRAIN, *Chrysichthys (filamentosus, nigrodigitatus, walkeri)* (visual sensory lobes in medulla) 264-266; *Hyla chrysoscelis* (seasonality of midbrain auditory responses) 844-852.

CALLS, *Hyla versicolor* (temp. effects on mating call) 733-738; *Physalaemus (coloradorum n. sp., petersi, pustulosus)* (advertisement calls compared) 902-921; *Rana yavapaiensis n. sp.* (call structure compared with spp. of *R. pipiens* complex) 940-949.

CARTILAGE, Ammodytidae, Pegasidae (enlarged upper jaw cartilages: functional & phylogenetic implications) 1011-1015.

CAUDAL GLAND, *Diplodactylus (ciliaris, rankini, strophurus)* (defensive secretion of) 1025-1028.

CHARACTERS, *Alosa pseudoharengus*, *Coregonus hoyi* (charc. displacement: evidence for competition?) 878-883; *Leiognathus elongatus* (2 spp. based on morph. & light organ charc.) 884-892; *Rana yavapaiensis n. sp.* (morphological & pattern charc. compared with spp. of *R. pipiens* complex) 940-949.

CHROMOSOMES, *Campostomus anomalum*, *Notemigonus crysoleucus*, *Notropis (emilia, lutrensis, venustus)*, *Pimephales vigilax* 133-139; Cyprinidae (cell culture technique for chromosome prep.) 232-235; *Gobionellus shufeldti* (multiple sex chromosomes) 235-238; *Ctenopharyngodon idella* × *Hypophthalmichthys nobilis* (chromosomal & electrophoretic analyses of) 337-342; *Eleutherodactylus (fitzingeri group, melanostictus, rugulosus group, unistrigatus series)* (karyology) 765-768; *Ictalurus (catus, furcatus, punctatus, & 4 hybrids of)* (karyology of) 873-878; *Rana yavapaiensis n. sp.* (karyotypes compared with spp. of *R. pipiens* complex) 940-949; *Noomis micropogon*, *Rhinichthys ("bowersi," cataractae)* (karyotypes) 990-992.

COLORATION, *Crotalus (atrox, molossus)* (melanism) 213-215; *Cichlasoma meeki* (color pattern variability & behav.) 534-538; *Sphaerodactylus micropithecus* (variation in color pattern) 768-770; *Grammistis sexlineatus* (coloration & probable toxicity in juv.) 1015-1017.

COMPETITION, *Anolis (cooki, cristatellus)* (competitive interference) 853-862; *Alosa pseudoharengus*, *Coregonus hoyi* (charc. displacement & habitat shift: competition evidence?) 878-883.

DEVELOPMENT, *Lagodon rhomboides*, *Diplodus holbrookii* (ontogenetic develop. & diet change) 174-187; *Ammodytes hexapterus* (egg, larvae) 242-244; *Thalassoma duperreyi* (develop. changes in reprod. tract during sex reversal) 245-248; *Poecilia latipinna* (seasonal maturation & growth in males) 252-255; *Lepisosteus* (cranial & pectoral girdle bones) 476-502; *Ichthyomyzon gagei* (metamorphosis of) 502-515; *Natrix tessellata* (embryonic devel. of Duvernoy's gland) 516-521; *Chelydra serpentina* (hydric environ. & embryonic metab.) 547-550; *Bulbonaricus davaensis* (metamorphosis from planktonic to demersal form) 565-571; *Litoria ewingi* (metamorphosis timing) 868-873.

DIGESTION, *Euthynmus lineatus* (gastric evacuation times) 1000-1005.

DISTRIBUTION, *Gasterosteus aculeatus* (in Boston, Mass.) 258-262; *Triportheus elongatus* (in Trinidad) 262-263; *Desmognathus fuscus* (heterozygosity & geogr. distr.) 343-356; *Engraulis (anchoita,*

edentulus, juriensis, mordax, mysticetus, ringens) (biogeogr., bipolar distr. of engraulids) 422-427; *Dicrolene intronigra* (depth dist.) 538-541; *Enchelycore anatina* (Mediterr. Israel) 541-544; *Myoconger compressus* (questionable distr. records) 585-594; *Platemys (radiolata, spixii)* (So. Amer. distr.) 780-786; *Pelamis platurus* (Galapagos records) 786-789.

ECOLOGY, *Stemonidium hypomelas* 249-252; *Esox lucius* (in an Alberta lake) 255-258.

ELECTROPHORESIS, *Bufo (americanus, fowleri)* 18-26; *Ictalurus (lupus, melas, punctatus)* 97-101; *Illyodon (furcidens, xantusi)* 102-107; *Gambusia affinis* 108-119; *Salmo (gairdneri, newberryi)* 120-132; *Hysterocarpus traski* 223-227; *Ctenopharyngodon idella* × *Hypophthalmichthys nobilis* (chromosomal & electrophoretic analyses of) 337-343; *Desmognathus fuscus* (heterozygosity) 343-356; *Rana (palmipes, punctulosa, tarahumarae, zweifeli n. sp.)* 398-403; *Notropis (chrysopoecalus, cornutus)* 617-628; *Stegastes fasciolatus* 629-640; *Rhinichthys bowersi* 652-659; *Eleutherodactylus (fitzingeri group, melanostictus, rugulosus group, unistrigatus series)* 765-768; *Gambusia puncticulata* complex 921-933; *Rana (a. aurora, aurora draytoni)* (electrophoretic differences) 1018-1022; *Diplodactylus (ciliaris, rankini, strophurus)* (electrophoretic analysis of defensive gland secretion) 1025-1028.

ENDANGERED SPECIES, *Gambusia amistadensis* (extinction of) 529-530.

EVOLUTION, kinosternid turtles (relationship to shell kinesis) 356-375; *Gasterosteus aculeatus* (body-size evol.) 530-534.

EXTINCTION, *Gambusia amistadensis* 529-530.

EYE, *Tantalophis discolor* (pupil) 215-216.

FECUNDITY, *Heterandria formosa* (fecundity variat. & body size) 720-726.

FEEDING, *Dermochelys coriacea* (benthic feeding) 552-553; *Crotalus horridus* 976-981; *Lepomis macrochirus* (group size & feeding rate) 998-1000; Amodytidae, Pegasidae (enlarged upper jaw cartilages: to enhance feeding function?) 1011-1015. FOAM, *Leptodactylus fuscus* (foam production in tadpoles) 778-780.

FOOD, *Nerodia rhombifera* (ontogenetic diet shift) 550-552; *Macrorhamphos scolopax* (diet & morphol. variat.: evidence for 2 spp.) 595-608; *Scaptochelys agassizii* 811-820; *Crotalus horridus* 976-981.

FOOTPRINTS, *Varanus komodensis* (analysis of footprint & locomotion; compar. with fossil footprints) 662-671.

FUNCTION, of teleost jaw protrusion 1-18.

GENETICS, *Gambusia affinis* (thermal, demographic effects on genetic differentiation) 108-119; *Salmo (gairdneri, newberryi)* 120-132; *Poecilia reticu-*

lata (heritability of dorsal ray no.) 140-144; *Ambystoma (platineum, texanum)* (gynogenesis) 324-337; *Desmognathus fuscus* (genetic variat.) 343-356; *Clupea harengus* (biochem. popn. genetics) 357-364; *Cyprinodon macularius* (evol. genetics) 364-369; *Stegastes fasciolatus* (genetic variat.) 629-640; *Etmopterus spinax* (quant. genetics of vertebrae no. & position of dorsal fin spines) 794-797; *Ictalurus (catus, furcatus, punctatus, & hybrids of)* (karyology of) 873-878.

GEOGRAPHIC LOCALITIES

Argentina, *Platemys spixii* 780-786.

Arizona, *Cyprinodon macularius* 364-369; *Rana yavapaiensis* n. sp. 940-948.

Arkansas, *Ctenopharyngodon idella* × *Hypophthalmichthys nobilis* 337-342; *Cnemidophorus sexlineatus* 826-832.

Atlantic (NW), *Dicrolene intronigra* 538-541; *Barathronus (multidens n. sp., unicolor n. sp.)* 579-584.

Australia, *Rhynchos (filamentosus, gloveri n. sp.)* 68-72; *Lumiconger arafura* 72-81; *Lates calcarifer* 144-149; *Synodus englemani* 187-194; *Echiodon rendahli* 410-422; *Dermochelys coriacea* 552-553; *Macrorhamphos scolopax* (2 spp.?) 595-608; *Litoria ewingi* 868-873; *Python s. spilotes* 959-965.

Austria, *Hyla arborea arborea* 609-616.

Alabama, *Ichthyomyzon gagei* 502-515; *Adinia xenica*, *Fundulus pulvereus* 797-800; *Ictalurus (catus, furcatus, punctatus)* 873-878.

Alaska, *Gasterosteus aculeatus* 530-534.

Bahama Is., *Gambusia (puncticulata manni, p. puncticulata)* 921-933.

Banda Sea, *Gorgasia barnesi* n. sp. 404-409.

Belize, *Gambusia puncticulata yucatana* 921-933.

Bermuda, *Fundulus (bermudae, n. sp.)* 820-825.

Bolivia, *Physalaemus petersi* 902-921.

Brazil, *Platemys (radiolata, spixii)* 780-786; *Physalaemus petersi* 902-921.

Canada, *Bufo (americanus × fowleri)* (Ont.) 18-26; *Micropterus dolomieu*, *Lepisosteus osseus* (Ont.) 149-152; *Coregonus clupeaformis* (Alb.) 238-242; *Esox lucius* (Alb.) 255-258; *Clupea harengus* (N. Bruns., N. Found.) 357-364.

California, *Hysterocarpus traski* 223-227; *Cyprinodon macularius* 364-369; *Uma inornata* 370-378; *Bufo boreas halophilus* 689-696; 696-701; *Hyla regilla* 727-732; *Scaptochelys agassizii* 811-820; *Rana muscosa* 966-976; *Rana (a. aurora, aurora draytoni)* 1018-1022.

Colombia, *Physalaemus (petersi, pustulosus)* 902-921.

Corsica, *Hyla arborea sarda* 609-616.

Costa Rica, *Eleutherodactylus melanostictus* 765-768.

Cuba, *Gambusia (puncticulata baracanna, puncticulata bucheri, puncticulata monticola, p. puncticulata)* 921-933.

Dakar, *Myroconger compressus* 585-594.

Denmark, *Clupea harengus* 357-364.

Ecuador, *Physalaemus (coloradorum n. sp., petersi)* 902-921.

Europe, *Gasterosteus aculeatus* 87-97.

Fiji, *Grammistes sexlineatus* 1015-1017.

- Florida, *Crocodylus acutus* 153–161; *Diplodus holbrooki*, *Lagodon rhomboides* 174–187; *Poecilia latipinna* 252–255; *Hyla crucifer* 302–308; *Heterantria formosa* 720–726; *Drymarchon couperi* 776–778; *Fundulus grandis* 820–825; *Hyla chrysoscelis* 844–852; *Eucinostomus (argenteus, gula, harengulus, jonesii)* 893–902; *Lepomis macrochirus* 996–998.
- France, *Hyla meridionalis* 609–616.
- Galapagos, *Pelamis platurus* 786–789; *Geochelone elephantopus vandenburghi* 702–707.
- Guatemala, *Plectrohyla (polomchi n. sp., tecunumani n. sp.)* 390–397.
- Grand Cayman I., *Gambusia p. puncticulata* 921–933.
- Hawaii, *Stemonidium hypomelas* 249–252; *Stegastes fasciatus* 629–640; *Labroides phthirophagus* 758–762; *Grammistes sexlineatus* 1015–1017.
- Iceland, *Anguilla* sp. 221–223.
- Idaho, *Salmo (gairdneri, newberryi)* 120–132.
- Illinois, *Ambystoma (platineum, platineum × texanum)* 324–337; *Desmognathus fuscus* 343–356.
- India, *Horaichthys setnai* 833–839.
- Indonesia, *Grammistes sexlineatus* 1015–1017.
- Inhaca I., *Echiodon cinereus* 410–422.
- Iowa, *Ictalurus punctatus* 873–878.
- Israel, *Mauremys caspica rivulata* 216–219; *Enchelycore anatina* 541–544; *Hyla savignyi* 609–616.
- Italy, *Hyla a. arborea* 609–616.
- Jamaica, *Anolis lineatopus* 747–757; *Gambusia p. puncticulata* 921–933.
- Japan, *Echiodon owasianus* 410–422; *Hyla japonica* 609–616; *Leiognathus (elongatus, stercorarius)* 884–892.
- Kansas, *Texasophis wilsoni* n. sp. 660–661; *Ictalurus punctatus* 873–878.
- Kentucky, *Desmognathus fuscus* 340–356; *Ambystoma texanum* 862–868.
- Korea, *Hyla (japonica, suweonensis)* 609–616.
- Louisiana, *Gobionellus shufeldti* 235–238; *Nerodia rhombifera* 550–552; *Akistodon piscivorus* 772–774; *Alligator mississippiensis* 839–843; *Hyla chrysoscelis* 844–852; *Ictalurus punctatus* 873–878.
- Maine, *Clupea harengus* 357–364.
- Marshall Is., *Grammistes sexlineatus* 1015–1017.
- Maryland, *Fundulus heteroclitus macrolepidotus* 820–825.
- Massachusetts, *Gasterosteus aculeatus* 258–262; *Desmognathus fuscus* 343–356; *Fundulus heteroclitus* 522–525; *Fundulus heteroclitus macrolepidotus* 820–825.
- Mauritius, *Grammistes sexlineatus* 1015–1017.
- Mexico, *Cyprinodon verecundus* n. sp. (Yuc.) 58–68; *Ilyodon (jal.)* 102–107; *Tantalophis discolor* (Oax.) 215–216; *Rana (pustulosa, trahumarae, zweifeli n. sp.)* 398–403; *Cichlasoma meeki* (Quin. Roo) 534–538; *Gambusia puncticulata yucatana* 921–933; *Cichlasoma socolofi* n. sp. 933–940.
- Michigan, *Desmognathus fuscus* 343–356.
- Mississippi, *Notropis umbratilis* 385–390.
- Missouri, *Hyla versicolor* 733–738; *Notropis nubilus* 994–996.
- Nebraska, *Chrysemys picta* 546–547; *Chelydra serpentina* 549–550.
- Nevada, *Gasterosteus doryssus* 530–534.
- New Jersey, *Desmognathus fuscus* 343–356.
- New Mexico, *Ictalurus (lupus, melas, punctatus)* 97–101; *Crotalus (atrox, molossus, viridis)* 213–215.
- New York, *Plethodon cinereus* 309–314.
- Nigeria, *Chrysichthys (filamentosus, nigrodigitatus, walkeri)* 264–266.
- North Carolina, *Desmognathus fuscus* 343–356; *Hyla chrysoscelis* 844–852.
- Norway, *Clupea harengus* 357–364.
- Ohio, *Desmognathus fuscus* 343–356; *Ictalurus punctatus* 873–878.
- Oklahoma, *Percina pantherina* 378–384.
- Oman, *Chelonla mydas* 774–776.
- Oregon, *Ambystoma gracile*, *Taricha granulosa* 672–683.
- Panama, *Eleutherodactylus melanostictus* 765–768.
- Pennsylvania, *Desmognathus fuscus* 343–356; *Crotalus horridus* 976–981.
- Peru, *Physalaemus petersi* 902–921.
- Philippines, *Genicanthus lamarchi* 194–200; *Echiodon anchipterus* n. sp. 410–422; *Leiognathus (elongatus, stercorarius)* 884–892.
- Puerto Rico, *Epipedes inornatus* 219–220; *Sphaerodactylus micropithecus* 768–770; *Anolis (cooki, cristatellus)* 853–862.
- Red Sea, *Grammistes sexlineatus* 1015–1017.
- Ryukyu Is., *Hyla hallowellii* 609–616.
- Singapore, *Gnatholepis hendersoni*, *Macrodonogobius wilburi* 227–229.
- Somalia, *Echiodon coheni* n. sp. 410–422.
- Solomon Is., *Emoia (rufilabialis, taumakoensis)* 571–578.
- South Carolina, *Gambusia affinis* 108–119; *Rana clamitans* 210–213; *Fundulus heteroclitus* 522–525; *Fundulus h. heteroclitus* 820–825; *Bufo terrestris* 1023–1025.
- Surinam, *Cnemidophorus lemniscatus* 713–719.
- Taiwan, *Hyla chinensis* 609–616.
- Tanzania, *Bulbonaricus brucei* 565–571.
- Tennessee, *Desmognathus fuscus* 343–356; *Etheostoma (Catonotus)* sp. 992–994.
- Texas, *Campostoma anomalam*, *Notemigonus crysoleucas*, *Notropis (emiliae, lutrensis, venustus)*, *Pimephales vigilax* 133–139; *Etheostoma spectabile* 291–301; *Eurycea (neotenes × tridentifera)* 428–441; *Menidia beryllina* 544–545; *Ictalurus punctatus* 873–878.
- Trinidad, *Triportheus elongatus* 262–263; *Leptodactylus fuscus* 778–780; *Physalaemus pustulosus* 902–921.
- Turkey, *Hyla savignyi* 609–616.
- Uruguay, *Platemys spixii* 780–786.
- Utah, *Gila (cypha, elegans)*, *Ptychocheilus lucius*, *Xyrauchen texanus* 790–793.
- Venezuela, *Hildebrandichthys setiger* 422–427; *Physalaemus pustulosus* 902–921.

- Vermont, *Desmognathus fuscus* 343-356.
 West Virginia, *Rhinichthys bowersi* 652-659.
 Wisconsin, *Ictalurus punctatus* 873-878.
 Zaire, *Nanochromis transversitus* 82-86.
- GROWTH**, *Cyprinodon (beltrani, labiosus, maya, simus, verecundus)* (growth/shape analysis) 58-68; *Ambystoma (jeffersonianum, maculatum)* (altitudinal effect on larval growth) 162-169; *Diplodus holbrooki*, *Lagodon rhomboides* (growth changes in teeth, mouth & body, & diet) 174-187; *Poecilia latipinna* (postmaturation growth in males) 252-255; *Anolis lineatopus* (seasonality of hatching recruitment & growth) 747-757; *Cyprinus carpio* (rel. growth of organs & parts; metab.-size relation) 800-803; *Ambystoma texanum* (larval growth) 862-868.
- HABITAT**, *Mauremys caspica rivulata* (in Israel) 216-219; *Esox lucius* (habitat utilization in an Alberta lake) 255-258; *Anolis* (development variabil. & habitat) 315-323; *Uma inornata* (wind-blown sand & habitat) 370-378; *Percina pantherina* (preferred habitat) 378-384; *Cyprinidae* (morphol. & habitat use) 442-455; *Desmognathus fuscus* (habitat moisture & activity) 684-688; *Bufo boreas hawaiiensis* (activity, microhabitat & thermal consequences) 689-696; *Natrix natrix* (movement, home range, habitat use) 707-713; *Cnemidophorus lemniscatus* (distr. & habitat of sexual & parthenogenetic popns.) 713-719; *Anolis (cooki, cristatellus)* (habitat & competition) 853-862; *Alosa pseudoharengus*, *Coregonus hoyi* (habitat shift) 878-883.
- HERMAPHRODITISM**, *Lates calcarifer* (protandry; sexually precocious popn.) 144-149; *Thalassoma duperreyi* (develop. changes in reprod. tract) 245-248; *Menidia beryllina* (spermatozoa & eggs in) 544-545.
- HISTOLOGY**, salamander spermatophore (microstructure & histochemistry) 739-747; *Horaichthys setnai* (testis structure) 333-339.
- HOME RANGE**, *Natrix natrix* (movements, home range, habitat use) 707-713.
- HYBRIDIZATION**, *Bufo (americanus × fowleri)* 18-26; *Ambystoma (platineum × texanum)* 324-337; *Ctenopharyngodon idella × Hypothalmichthys nobilis* (chromosomal & electrophoretic analyses of) 337-342; *Eurycea (neotenes × tridentifera)* 428-441; *Hyla (a. arborea, arborea sarda, chinensis, hallowellii, japonica, meridionalis, savignyi, suweonensis)* (systematic implications of hybridization experiments) 609-616; *Rana yavapaiensis* n. sp. (experimental hybridization with 4 other spp. of *Rana pipiens* complex) 940-949; *Notropis (chrysoccephalus, cornutus)* 617-628; *Ictalurus (punctatus × furcatus, furcatus × punctatus, punctatus × catus, catus × furcatus)* (karyotypes of) 873-878.
- IMMUNOLOGY**, *Heterodon*, *Farancia*, *Carphophis*, *Conophis*, *Diadophis*, *Contia* (immunological studies of neotropical xenodontine snakes) 641-652.
- JAWS**, *Ammodytidae*, *Pegasidae* (enlarged upper jaw cartilages: phylogenetic & functional implications) 1011-1015.
- KIDNEY**, *Salmo gairdneri* (distr. of renal corpuscles) 264.
- KINESIS**, shell kinesis in kinosternid turtles 456-475.
- LARVAE**, *Ambystoma (jeffersonianum, maculatum)* 161-169; *Ammodytes hexapterus* 242-244; *Ichthyomyzon gagei* (metamorphosis of ammocoetes) 502-515; *Leptodactylus fuscus* (foam production in tadpoles) 778-780; *Ambystoma texanum* (growth & survivorship) 862-868; *Physalaemus (coloradorum n. sp., petersi, pustulosus)* (larvae compared) 902-921; *Paralichthys lethostigma* (salinity, formalin conc. & buffer on larvae preservation) 981-988.
- LIFE HISTORY**, *Ambystoma (jeffersonianum, maculatum)* (altitudinal pattern of larval life history traits) 161-169; *Notropis umbratilis* (in Miss.) 385-390.
- LIGHT**, *Bufo boreas* (nocturnal activity & ambient illumination) 283-290.
- LIGHT ORGAN**, *Leiognathus (elongatus, stercorarius)* (2 spp. based on) 884-892.
- LOCOMOTION**, *Varanus komodensis* (analysis of footprints & locomotion; comparison with fossil footprints) 662-671; *Alligator mississippiensis* (kinematics of undulatory swimming) 839-843; *Euthynnus lineatus* (swimming performance) 1000-1005.
- LUMINESCENCE**, in *Lumiconger arafura* 72-81; *Leiognathus (elongatus, stercorarius)* (2 spp. based on light-organ charc.) 884-892.
- METABOLISM**, *Chelydra serpentina* (hydric environ. & embryonic metab.) 547-550; *Cyprinus carpio* (rel. growth with ref. to metab.-size relationship) 800-803.
- METHODS**, nerve staining (Sudan Black) 204-208; blood sampling from turtles 209-210; preparing disarticulated small fish skeletons 230-232; cell culture technique for chromosome prep. 232-235; ichthyoplankton preservation in formalin 525-528; *Paralichthys lethostigma* (salinity, formalin conc. & buffer effects on quality of larval preservation) 981-988; alizarin fish staining without causing exfoliation of scales 989-990.
- MORPHOLOGY**, of jaw protrusion of teleosts 1-18; *Lagodon rhomboides*, *Diplodus holbrooki* (ontogenetic & interspecific differ. in diet & morph.) 174-187; *Anolis* (variation in develop. morph.) 315-323; *Cyprinidae* (morph. & habitat use) 442-455; *Kinosternidae* (morph. design & shell kinesis) 356-375; salamander spermatophores (microstructure of) 739-747; *Cnemidophorus sexlineatus* (egg-shell morph.) 826-832; *Horaichthys setnai* (testis structure) 833-839; *Eucinostomus (argenteus, gula, harengulus, jonesi)* (morph. analysis) 893-902.

MOVEMENT, *Natrix natrix* (seasonal movements, home range, habitat use) 707–713; *Phyllodactylus lanei* (tail autotomy inhibits tonic immobility) 763–764.

MUSCLES, *Eleutherodactylus (fitzingeri group, melanostictus, rugulosus group, unistrigatus series)* (jaw muscle charc.) 765–768.

NERVES, nerve stain (Sudan Black B) 204–208.

NESTING, *Crocodylus acutus* (nest environ.) 153–161; *Geochelone elephantopus vanderburghi* (impact of feral burros on nest success) 702–707; *Python s. spilotes* (temp. elevation by shivering by brooding female) 959–965; *Etheostoma (Catonotus) sp.* (an egg-clusterer) 992–994.

OSTEOLOGY, of *Lumiconger arafura* 72–81; preparation of disarticulated fish skeletons 230–232; *Lepisosteus* (cranial & pectoral girdle bones) 476–502; *Myroconger compressus* 585–594; “*Morone saxatilis*” skull in Gregory’s “Fish Skulls” an *Epinephelus* (probably *striatus*) 793; *Physalaemus (coloradorum n. sp., petersi, pustulosus)* (compared) 902–921; *Pterygoplichthys* sp. (strength of pectoral girdle/spine complex) 1005–1008.

PALEONTOLOGY, *Gasterosteus dorissus* (Mio., Nev.) 530–534; *Gasterosteus aculeatus* (Rec., Alaska) 530–534; *Texasophis wilsoni* n. sp. (Mio., Kansas) 660–661; analysis of fossil footprints 662–672.

PECTORAL GIRDLE, *Pterygoplichthys* sp. (pectoral spine/girdle strength) 1005–1008.

PHYSIOLOGY, *Python s. spilotes* (shivering thermogenesis) 959–965; *Rana muscosa* (behavioral temp. modulation) 966–976; *Euthynnis lineatus* (gastric evacuation times) 1000–1005.

POPULATION, *Gambusia affinis* (microgeogr. popns. in S. Carolina reservoir) 108–119; *Lates calcarifer* (sexually precocious popn.) 144–149; *Hysterocephalus traski* (electrophoretic var. in popns.) 222–227; *Gasterosteus aculeatus* (low & complete morphs in Boston, Mass.) 258–262; *Desmognathus fuscus* (heterozygosity & geogr. distr. of popns. of) 343–356; *Clupea harengus* (biochem. popn. genetics) 357–364; *Cyprinodon macularius* (evol. genetics of) 364–369; *Notropis umbratilis* (popn. age structure) 385–390; *Eurycea (neotenes, tridentifera)* (popn. var.) 428–441; *Sitostoma fasciatum* (popn. structure in Hawaiian Archip.) 629–640; *Ambystoma gracile*, *Taricha granulosa* (high-density & low-density popn.) 672–683; *Cnemidophorus lemniscatus* (distr. & habitat of sexual & parthenogenetic popns. in Surinam) 713–719; *Cottus (k. klamathensis, k. polyporus n. subsp.)* (geogr. popns. compared) 949–954.

PREDATION, *Synodus englemani* (predator/prey relationship; predatory behav.) 187–194; *Epicrates*

inornatus (predation on bats) 219–220; *Nerodia rhombifera* (ontogenetic diet shift) 550–552. **PRESERVATION**, *Paralichthys lethostigma* (larvae preservation quality & effects of salinity, formalin conc. & buffer) 981–988.

RADIO-TELEMETRY, *Natrix natrix* (to determine home range, movement & habitat use) 707–713.

RECRUITMENT, *Anolis lineatopus* (hatching recruitment & growth) 747–757.

REPRODUCTION, *Poeciliopsis lucida* (male mating selectivity) 169–174; *Genicanthus lamarck* (reprod. behav.) 194–200; *Coregonus clupeaformis* (fecundity variation in Alberta) 238–242; *Ammodytes hexapterus* (lab. spawning) 242–244; *Poecilia latipinna* (seasonal maturation) 252–255; *Hyla crucifer* (egg-size variat.) 302–308; *Ambystoma (platineum, texanum)* (gynogenesis) 324–337; *Notropis umbratilis* (reprod. periodicity, fecundity) 385–390; *Fundulus heteroclitus* (variat. in spawning site) 522–525; *Dicrolene intronigra* (spawning season, maturity, fecundity) 538–541; *Notropis (chryscephalus, cornutus)* (level of reprod. isolation) 617–628; *Geochelone elephantopus vanderburghi* (hatching success & feral burro trampling) 702–707; *Heterandria formosa* (body size & fecundity) 720–726; *Hyla regilla* (mating behav. success) 727–732; *Adinia xenica*, *Fundulus pulvereus* (spawning associated with semilunar tidal cycles) 797–800; *Scaphiophryne agassizii* 811–820; *Fundulus (bermudae, grandis, h. heteroclitus, heteroclitus macrolepidotus, n. sp.)* (reprod. behav.) 820–825; *Horaichthys setnai* (testis structure & formation of spermatophores) 833–839; *Etheostoma (Catonotus) sp.* (courtship behav., nests, nesting) 992–994; *Notropis nubilus* (spawning activity & eggs) 994–996; *Bufo terrestris* (amplexus displacement) 1023–1025.

SCALES, *Lepisosteus* (development) 476–502; *Sphenodon punctatus micropithecus* (variat. in scale counts) 768–770; *Drymarchon corais couperi* (sex dimorph. in keeled scales) 776–778.

SEASONALITY, *Etheostoma spectabile* (seasonal variat. in egg size) 291–301; *Anolis* (dewlap variability & seasonality) 315–323; *Notropis umbratilis* (reprod. periodicity) 385–390; *Bufo boreas halophilus* (seasonal activity) 689–696; *Natrix natrix* (seasonal movement) 707–713; *Anolis lineatopus* (seasonality of hatching recruitment & growth) 747–757; *Hyla chrysoscelis* (seasonality of midbrain auditory responses) 844–852.

SEX, *Lates calcarifer* (sexually precocious popn.) 144–149; *Thalassoma duperrey* (sex reversal & anatomical changes) 245–248; *Chrysemys picta* (sex differentiation/threshold temp.) 545–547; *Cnemidophorus lemniscatus* (distr. & habitats of sexual & parthenogenetic popns. in Surinam) 713–719; *Chelonia mydas* (sex ratio in Oman) 774–776.

SEXUAL DIMORPHISM. *Nanochirus transvestitus* n. sp. 82–86; *Stemonidium hypomelas* 249–252; *Macrorhamphosus scolopax* (diet & morph. evidence for 2 sexually dimorph. spp.) 595–608; *Drymarchon corais couperi* (sex. dimorph. in keeled dorsal scales) 776–778.

SIZE. *Gasterosteus aculeatus* (gigantism; body size evol.) 530–534; *Dicrolene introna* (size-depth relat.) 538–541; *Ambystoma gracile*, *Taricha granulosa* (size & competition) 672–678; *Heterandria formosa* (body size & fecundity) 720–726; *Cyprinus carpio* (rel. growth with ref. to metab.-size relat.) 800–803; *Rana (a. aurora, aurora draytoni)* (size differences) 1018–1022.

SPERMATOPORES, microstructure & histochem. of salamander spermatophores 739–747.

SPINES, *Etmopterus spinax* (quant. genetics of vertebrae no. & dorsal spine position) 794–797.

SURVIVAL, *Scaptochelys agassizii* (in Inyanpah Valley, Calif.) 811–820; *Ambystoma texanum* (embryonic & larval survivorship) 862–868.

SYMBIOSIS, *Labroides phthirophagus* (preference for ectoparasite-infected host fish) 758–762; *Lepomis macrochirus*, *Mugil cephalus* (cleaning by bluegill in Fla.) 996–998.

SYSTEMATICS, *Allenichthys* n. gen., *Antennatus*, *Echinophryne*, *Histiophryne*, *Histrio*, *Kuiterichthys* n. gen., *Lophiocharon*, *Nudiantennarius*, *Phyllophryne* n. gen., *Rhycherus*, *Tathicarpus*, *Trichophryne* (generic revision of Antennariidae) 27–44; *Bathyraja*, *Gurgesiella*, *Neoraja*, *Pavoraja*, *Raja*, tribe *Pavorajini* nov. (gen. revision) 45–58; *Cyprinodon verecundus* n. sp. (Laguna Chichancanab, Yuc., Mex.) 58–68; *Rhycherus (bifurcatus, filamentosus, gloveri* n. sp. [Australia], *wildii*) 68–72; *Lumiconger arafura* n. sp. (n. Australia) 72–81; *Nanochromis transvestitus* n. sp. (Zaire) 82–86; *Rana clamitans* (redescr. of holotype) 210–213; *Rivulichthys tuerlingi* (synon. of *Pterolebias longipinnis*) 528–529; *Italurus (lupus, melas, punctatus)* (electrophoretic analysis) 97–101; *Polyodon spathula* (neglected account of) 201–204; *Gnatholepis hendersoni*, *Macrodonogobius wilburi* (synonymization) 227–229; *Plectrohyla (pokomchi* n. sp., *tecuumanii* n. sp.) (Guatemala) 390–397; *Rana (tarahumarae, zweifeli* n. sp. [sw. Mex.]) *pustulosa* 398–403; *Gorgasia barnesi* n. sp. (Banda Sea) 404–409; *Echiodon (anchipterus* n. sp. [Philip.], *coheni* n. sp. [Somalia], *cinereus*, *ouwasianus*, *rendahli*) 410–422; *Engraulis (subgen. Cetengraulis) (anchoita, edentulus, juruensis, mordax, mysticetus, ringens)* (revision) 422–427; *Bulbonaricus* (sr. synon. *Enchelyscampus*), *B. brucei* n. sp. (e. Afr.) 565–571; *Emoia (caeruleoocauda, cyanura, schmidti, maculata, rufilabialis* n. sp. [Solomon Is.], *taumakoensis* n. sp. [Solomon Is.]) 571–578; *Barathronus (multidens* n. sp., *unicolor* n. sp.) (nw. Atlan.) 579–584; *Myroconger compressus* (redescr.) 585–594; *Hyla (a. arborea, arborea sarda, chinensis, hallowellii, japonica, meridionalis, savignyi, suweonensis)* (system. implications of hybrid-

ization experiments) 609–616; *Heterodon*, *Fafrancia*, *Carphophis*, *Conophis*, *Diadophis*, *Contia* (molecular systematics of neotropical xenodontine snakes) 641–652; *Rhinichthys boweri* (a valid species, not a hybrid) 652–659; *Texasophis wilsoni* n. sp. (Mio., Kansas) 660–661; *Eleutherodactylus melanostictus* (phylogenetic relationships) 765–768; “*Morone saxatilis*” skull in Gregory’s “Fish Skulls” a skull of *Epinephelus* (probably *striatus*) 793–794; *Leiognathus (elongatus, stercorarius)* (2 spp. based on morphological & light organ charc.) 884–892; *Eucinostomus (argenteus, gula, harengulus, jonesi)* (morphol. analysis, recognition of 4 spp. in *E. argenteus* complex) 893–902; *Physalaemus (coloradorum* n. sp. [Ecuador], *petersi*, *pustulosus*) 902–921; *Gambusia (puncticulata complex, yucatana complex)* (taxonomically distinguished) 921–933; *Cichlasoma socolofi* n. sp. (Chia., Mex.) 933–940; *Rana yavapaiensis* n. sp. (Ariz.) 940–948; *Cottus (klamathensis polyporus* n. subsp. [Calif.], *k. klamathensis*, *k. macrops*) (popns. examined) 949–957; *Phycis musicki* (new name to replace *Phycis tenuis* [Koken, 1891], a homonym of *Phycis tenuis* [Mitchell, 1814]) 1008–1009; *Uranoscopus kaianus* (redescr. & lectotype designation) 1009–1011; *Ammodytidae*, *Pegasidae* (phylogenetic implications of enlarged upper jaw cartilages) 1011–1015.

TEMPERATURE, *Gambusia affinis* (genetic diff. & thermal conditions) 108–119; *Chrysemys picta* (low threshold temp. for sex determination) 546–547; *Bufo boreas halophilus* (activity pattern & thermal biol.) 689–696; (temp. & daily emergence time) 696–701; *Hyla versicolor* (temp. effects on mating call) 733–738; *Python s. spilotes* (body temp. elev. by shivering in brooding female) 959–965; *Rana muscosa* (temp. modulation in) 966–976; *Euthynus lineatus* (swimming body temp.) 1000–1005.

TOXICITY, *Grammistes sexlineatus* (juvs. toxic?) 1015–1017.

VARIATION, *Gasterosteus aculeatus* (geogr. variat.) 87–97; *Ilyodon (furcidens/xantusi)* (geogr. variat.) 102–107; *Campostomus anomalum*, *Notemigonus crysoleucus*, *Notropis (emiliae, lutrensis, venustus)*, *Pimephales vigilis* (chromosomal variat.) 133–139; *Diplodus holbrookii*, *Lagodon rhomboides* (ontogenetic & interspecific variat. in diet) 174–187; *Hysterocarpus traski* (electrophoretic variat.) 222–227; *Coregonus clupeaformis* (fecundity variat.) 238–242; *Etheostoma spectabile* (egg size variat.) 291–301; *Hyla crucifer* (intraclutch egg size variat.) 302–308; *Anolis* (interspecific variat. of dewlap) 315–323; *Desmognathus fuscus* (genetic variat.) 343–356; *Eurycea (neotenes, tridentifera)* (popn. variat.) 428–441; *Fundulus heteroclitus* (variats. in spawning sites) 522–525; *Macrorhamphosus scolopax* (diet/ morphol. variat.) 595–608; *Stegastes fasciatus*

(genetic variat.) 629-640; *Heterandria formosa* (fecundity variat. & body size) 720-726; *Sphaerodactylus micropithecus* (variats. in dorsal pattern & scale counts) 768-770; *Cottus klamathensis* (geogr. variat.) 949-959.

VERTEBRAE, *Etmopterus spinax* (quant. genetics of vertebrae no. & position of dorsal spines) 794-797.